

To appear in *Algebra Colloquium*

SQUAREFREE MONOMIAL MODULES AND EXTREMAL BETTI NUMBERS

MARILENA CRUPI AND CARMELA FERRÒ

University of Messina, Department of Mathematics and Computer Science
Viale Ferdinando Stagno d'Alcontres, 31
98166 Messina, Italy
E-mail: mcrupi@unime.it; cferro@unime.it

ABSTRACT. Let K be a field and let $S = K[x_1, \dots, x_n]$ be a polynomial ring over K . Let $F = \oplus_{i=1}^r S e_i$ be a finitely generated graded free S -module with basis $e_1, \dots, e_r$ in degrees $f_1, \dots, f_r$ such that $f_1 \leq f_2 \leq \dots \leq f_r$. We examine some classes of squarefree monomial submodules of F . Hence, we focalize our attention on the Betti table of such classes in order to analyze the behavior of their extremal Betti numbers.

2010 Mathematics Subject Classification: 05E40, 13B25, 13D02 16W50.

Keywords: graded modules, squarefree monomial modules, minimal graded resolutions.

INTRODUCTION

Let K be a field and let $S = K[x_1, \dots, x_n]$ be the polynomial ring in n variables with coefficients in K . Let $F = \oplus_{i=1}^r S e_i$ be a finitely generated graded free S -module with basis $e_1, \dots, e_r$ in degrees $f_1, \dots, f_r$, renumbered as necessary so that $f_1 \leq f_2 \leq \dots \leq f_r$.

A *monomial submodule* M of F is a submodule generated by monomials, *i.e.*, $M = \oplus_{i=1}^r I_i e_i$, where I_i are the monomial ideals of S generated by those monomials m of S such that $m e_i \in M$. A monomial submodule $M = \oplus_{i=1}^r I_i e_i \subsetneq F$ is a *squarefree monomial submodule* if every I_i is a squarefree monomial ideal of S .

If $F = S$, then a squarefree monomial submodule of F is a squarefree monomial ideal of S . The class of squarefree monomial ideals has been studied from viewpoint of commutative algebra and combinatorics (see, for example [1, 2]).

In this paper, we are interested in the study of some classes of squarefree monomial submodules: *squarefree stable submodules*, *squarefree strongly stable submodules* and *squarefree lexicographic submodules*.

As in the theory of squarefree monomial ideals, we have the following hierarchy of squarefree monomial submodules: squarefree lexicographic submodule $\Rightarrow$ squarefree strongly stable submodule $\Rightarrow$ squarefree stable submodule.

In [1], Aramova, Herzog and Hibi constructed the explicit minimal graded free resolution of a squarefree stable ideal, similar to the Eliahou-Kervaire resolution of a stable ideal [11], and stated a formula for computing the Betti numbers of such class of ideals. Such formula is a fundamental tool if one wants to study the Betti table of a squarefree stable submodule.

If M is a finitely generated graded S -module, a Betti number $\beta_{k,k+\ell}(M) \neq 0$ is called *extremal* if $\beta_{i,i+j}(M) = 0$ for all $i \geq k$, $j \geq \ell$, $(i, j) \neq (k, \ell)$.

The extremal Betti numbers were introduced by Bayer, Charalambous and Popescu in [3] as a refinement of two invariants of the module M : the projective dimension and the Castelnuovo-Mumford regularity.

Indeed, if $\beta_{k_1, k_1+\ell_1}(M), \dots, \beta_{k_t, k_t+\ell_t}(M)$, $k_1 > \dots > k_t, \ell_1 < \dots < \ell_t$, are all extremal Betti numbers of M , then $k_1 = \text{projdim}_S(M)$ and $\ell_t = \text{reg}_S(M)$.

The behavior of the extremal Betti numbers for some classes of monomial ideals in polynomial rings in a finite numbers of variables over a field was studied by Crupi and Utano in [7, 8]. Subsequently, the same authors [9] examined such special graded Betti numbers for a lexicographic submodule M of a finitely generated graded free S -module.

In this paper we devote our attention on the extremal Betti numbers of a square-free stable submodule. We characterize the extremal Betti numbers of such class of monomial submodules and give a criterion for determining their positions and their number.

The plan of the paper is the following. In Section 1, some notions that will be used throughout the paper are recalled. In Section 2, the classes of squarefree stable submodules, squarefree strongly stable submodules and squarefree lexicographic-submodules are examined and some relevant properties discussed. In Section 3, the behavior of the extremal Betti numbers for the squarefree stable submodules is studied; the characterization of these invariants is the main result. In Section 4, given a squarefree stable submodule M of the free S -module S^r , $r \geq 1$, a criterion to recognize among all the graded Betti numbers of M those extremal ones is given. Section 5 contains an application on the so called *super extremal Betti numbers* of a squarefree lexicographic submodule.

1. PRELIMINARY AND NOTATION

Throughout this paper, let $S = K[x_1, \dots, x_n]$ be the polynomial ring in n variables over a field K and $F = \bigoplus_{i=1}^r S e_i$ a finitely generated graded free S -module with basis $e_1, \dots, e_r$ in degrees $f_1, \dots, f_r$ such that $f_1 \leq f_2 \leq \dots \leq f_r$. We consider S as an $\mathbb{N}$ -graded ring and each $\deg x_i = 1$.

The elements of the form $x^a e_i$, where $x^a = x_1^{a_1} x_2^{a_2} \dots x_n^{a_n}$ for $a = (a_1, \dots, a_n) \in \mathbb{N}_0^n$, are called *monomials* of F .

A monomial $m e_i \in F$ is called a *squarefree monomial* of F if m is a squarefree monomial of S , i.e., $m = x_{i_1} x_{i_2} \dots x_{i_d}$ with $1 \leq i_1 < i_2 < \dots < i_d \leq n$.

For every monomial $m e_i \in F$, we set

$$\deg(m e_i) = \deg(m) + \deg(e_i).$$

In particular if $F \simeq S^n$ and $e_i = (0, \dots, 0, 1, 0, \dots, 0)$, where 1 appears in the i -th place, one has

$$\deg(m e_i) = \deg(m).$$

A *monomial submodule* M of F is a submodule generated by monomials, *i.e.*, $M = \oplus_{i=1}^r I_i e_i$, where I_i are the monomial ideals of S generated by those monomials m of S such that $me_i \in M$ [12].

A monomial submodule $M = \oplus_{i=1}^r I_i e_i \subsetneq F$ is a *squarefree monomial submodule* if every I_i is a squarefree monomial ideal of S , *i.e.*, every I_i is a monomial ideal of S generated by squarefree monomials.

For every monomial ideal $I \subsetneq S$, we denote by $G(I)$ the unique minimal set of monomial generators of I , by $G(I)_\ell$ the set of monomials v of $G(I)$ such that $\deg v = \ell$ and by $G(I)_{>\ell}$ the set of monomials u of $G(I)$ such that $\deg u > \ell$.

For every monomial submodule $M = \oplus_{i=1}^r I_i e_i$ of F such that $I_i \subsetneq S$, for $i = 1, \dots, r$, we set

$$\begin{aligned} G(M) &= \{ue_i : u \in G(I_i), i = 1, \dots, r\}, \\ G(M)_\ell &= \{ue_i : u \in G(I_i)_{\ell-f_i}, i = 1, \dots, r\}, \\ G(M)_{>\ell} &= \{ue_i \in G(M) : u \in G(I_i)_{>\ell-f_i}, i = 1, \dots, r\}. \end{aligned}$$

Let M be a finitely generated graded S -module, then M has a minimal graded free S -resolution

$$F : 0 \rightarrow F_s \rightarrow \dots \rightarrow F_1 \rightarrow F_0 \rightarrow M \rightarrow 0,$$

where $F_i = \oplus_{j \in \mathbb{Z}} S(-j)^{\beta_{i,j}}$. The integers $\beta_{i,j} = \beta_{i,j}(M) = \dim_K \text{Tor}_i(K, M)_j$ are called the graded Betti numbers of M .

Definition 1.1. A Betti number $\beta_{k,k+\ell}(M) \neq 0$ is called *extremal* if $\beta_{i,i+j}(M) = 0$ for all $i \geq k$, $j \geq \ell$, $(i, j) \neq (k, \ell)$.

The pair (k, ℓ) is called a *corner*.

2. SQUAREFREE MONOMIAL SUBMODULES

In this Section, we analyze some classes of squarefree monomial submodules of the finitely generated graded free S -module $F = \oplus_{i=1}^r S e_i$.

If I is a graded ideal of the polynomial ring S , we denote by $\text{indeg} I$ the *initial degree* of I , *i.e.*, the minimum t such that $I_t \neq 0$.

For a monomial $1 \neq u \in S$, we set

$$\text{supp}(u) = \{i : x_i \text{ divides } u\},$$

and

$$\text{m}(u) = \max\{i : i \in \text{supp}(u)\}.$$

Moreover, we set $\text{m}(1) = 0$.

Definition 2.1. Let $I \subsetneq S$ be a squarefree monomial ideal. I is called a *squarefree stable ideal* if for all $u \in G(I)$ one has $(x_j u)/x_{\text{m}(u)} \in I$ for all $j < \text{m}(u)$, $j \notin \text{supp}(u)$. I is called a *squarefree strongly stable ideal* if for all $u \in G(I)$ one has $(x_j u)/x_i \in I$ for all $i \in \text{supp}(u)$ and all $j < i$, $j \notin \text{supp}(u)$.

For every $1 \leq d \leq n$, let $[x_1, \dots, x_n]^d$ be the squarefree monomial ideal of S whose minimal system of monomial generators is given by all the degree d squarefree monomials in the variables $x_1, \dots, x_n$.

For example, for $n = 4$ and $d = 3$,

$$[x_1, \dots, x_4]^3 = (x_1x_2x_3, x_1x_2x_4, x_1x_3x_4, x_2x_3x_4).$$

Following [14], we give the following definition.

Definition 2.2. A submodule M of F is a *squarefree stable submodule* if

- (1) $M = \oplus_{i=1}^r I_i e_i$ is a squarefree monomial submodule with $I_i \subsetneq S$, for $i = 1, \dots, r$;
- (2) for every squarefree monomial $ue_i \in M$, $u \in S$, then $x_j \frac{u}{x_{m(u)}} e_i \in M$, for all $j < m(u)$, $j \notin \text{supp}(u)$;
- (3) $[x_1, \dots, x_n]^{f_j - f_i} I_j \subseteq I_i$ for every $j > i$.

Proposition 2.3. Let $M \subseteq F$ be a graded submodule.

Then M is a squarefree stable submodule of F if and only if

- (i) $M = \oplus_{i=1}^r I_i e_i$, with $I_i \subsetneq S$ squarefree stable ideal, for $i = 1, \dots, r$, and
- (ii) $[x_1, \dots, x_n]^{f_{i+1} - f_i} I_{i+1} \subseteq I_i$, for $i = 1, \dots, r - 1$.

Proof. In order to prove the assert it is sufficient to show that condition (3) in Definition 2.2 is equivalent to the following inclusion:

$$(2.1) \quad [x_1, \dots, x_n]^{f_{i+1} - f_i} I_{i+1} \subseteq I_i,$$

for $i = 1, \dots, r - 1$.

It is clear that statement (2.1) follows from condition (3) for $j = i + 1$.

Hence, in order to prove the required equivalence, it is sufficient to verify that

$$[x_1, \dots, x_n]^{f_{i+t} - f_i} I_{i+t} \subseteq I_i, \quad \text{for } t \geq 1.$$

We proceed by induction on t .

For $t = 1$ the assert follows from (2.1).

Now take $t > 1$ and suppose $[x_1, \dots, x_n]^{f_{i+t} - f_i} I_{i+t} \subseteq I_i$.

Since

$$[x_1, \dots, x_n]^{f_{i+t+1} - f_i} I_{i+t+1} = [x_1, \dots, x_n]^{f_{i+t+1} - f_{i+t} + f_{i+t} - f_i} I_{i+t+1},$$

from (2.1), we have

$$[x_1, \dots, x_n]^{f_{i+t+1} - f_i} I_{i+t+1} \subseteq [x_1, \dots, x_n]^{f_{i+t} - f_i} I_{i+t},$$

and from the inductive hypothesis we get the stated result. $\square$

Furthermore, we give the following definition.

Definition 2.4. A submodule M of F is a *squarefree strongly stable submodule* if

- (1) $M = \oplus_{i=1}^r I_i e_i$, with $I_i \subsetneq S$ squarefree strongly stable ideal, for $i = 1, \dots, r$;
- (2) $[x_1, \dots, x_n]^{f_{i+1} - f_i} I_{i+1} \subseteq I_i$, for $i = 1, \dots, r - 1$.

Remark 2.5. If M is a graded submodule of the finitely generated graded free S -module S^r , then M is a squarefree (strongly) stable submodule of S^r if and only if $M = \oplus_{i=1}^r I_i e_i$, with $I_i \subsetneq S$ squarefree (strongly) stable ideal, for $i = 1, \dots, r$, and

$$I_r \subseteq I_{r-1} \subseteq \dots \subseteq I_1.$$

For every $1 \leq d \leq n$, let $\{x_1, \dots, x_n\}^d$ be the set of all squarefree monomials of degree d in the variables $x_1, \dots, x_n$. We can order $\{x_1, \dots, x_n\}^d$ with the *squarefree lexicographic order* $\geq_{\text{slex}}$ [1].

Precisely, let

$$u = x_{i_1}x_{i_2}\cdots x_{i_d}, \quad v = x_{j_1}x_{j_2}\cdots x_{j_d},$$

with $1 \leq i_1 < i_2 < \cdots < i_d \leq n$, $1 \leq j_1 < j_2 < \cdots < j_d \leq n$, be squarefree monomials of degree d in S , then

$$u >_{\text{slex}} v \quad \text{if} \quad i_1 = j_1, \dots, i_{s-1} = j_{s-1} \quad \text{and} \quad i_s < j_s,$$

for some $1 \leq s \leq d$.

A nonempty set $L \subseteq \{x_1, \dots, x_n\}^d$ is called a *squarefree lexsegment set* of degree d if for $u \in L$, $v \in \{x_1, \dots, x_n\}^d$ such that $v >_{\text{slex}} u$, then $v \in L$.

Definition 2.6. Let $I \subsetneq S$ be a graded ideal. I is a *squarefree lexsegment ideal of degree d* if I is generated by the squarefree monomials belonging to a squarefree lexsegment set of degree d .

I is a *squarefree lexsegment ideal* if for all $1 \leq d \leq n$, if $u, v \in \{x_1, \dots, x_n\}^d$ with $u \in I$ and $v >_{\text{slex}} u$, then $v \in I$.

For every $d \geq 1$, let F_d be the part of degree d of $F \oplus_{i=1}^r Se_i$. Denote by $M^s(F_d)$ the set of all squarefree monomials of degree d of F . We order such set using the ordering $>_{\text{slex}}$, above defined. We will denote the new ordering by $>_{\text{slex}_F}$.

It is defined as follows: if ue_i and ve_j are squarefree monomials of F such that $\deg(ue_i) = \deg(ve_j)$, then

$$ue_i >_{\text{slex}_F} ve_j \quad \text{if} \quad \begin{cases} i < j \text{ or} \\ i = j \text{ and } u >_{\text{slex}} v. \end{cases}$$

Definition 2.7. Let $\mathcal{L} = \oplus_{i=1}^r I_i e_i$ be a squarefree monomial submodule of F such that $I_i \subsetneq S$, for $i = 1, \dots, r$. We call $\mathcal{L}$ a *squarefree lexicographic submodule* if for each degree $d \geq 1$, if $u, v \in M^s(\mathcal{L}_d)$ with $u \in \mathcal{L}$ and $v >_{\text{slex}_F} u$, then $v \in \mathcal{L}$.

Proposition 2.8. Let $\mathcal{L} \subsetneq S$ be a graded submodule.

Then $\mathcal{L}$ is a squarefree lexicographic submodule of F if and only if

- (i) $\mathcal{L} = \oplus_{i=1}^r I_i e_i$, with $I_i \subsetneq S$ squarefree lexsegment ideal, for $i = 1, \dots, r$, and
- (ii) $[x_1, \dots, x_n]^{\rho_i + f_i - f_{i-1}} \subseteq I_{i-1}$, for $i = 2, \dots, r$, with $\rho_i = \text{indeg } I_i$.

Proof. Let $\mathcal{L}$ be a squarefree lexicographic submodule of F .

(i) Since $\mathcal{L}$ is a squarefree monomial submodule of F , one has $\mathcal{L} = \oplus_{i=1}^r I_i e_i$, with I_i squarefree monomial ideal of S , for every i . Let $u, v \in \{x_1, \dots, x_n\}^d$ with $u \in I_i$ and $v >_{\text{slex}} u$. It follows that $ve_i >_{\text{slex}_F} ue_i$. Since $ue_i \in I_i e_i$ and $\mathcal{L}$ is a squarefree lexicographic submodule of F , $ve_i \in I_i e_i$, and so $v \in I_i$, i.e., I_i is a squarefree lexsegment ideal of S for every i .

(ii) Since I_i is a squarefree lexsegment ideal of S , then $x_1 x_2 \cdots x_{\rho_i} \in I_i$, $\rho_i = \text{indeg } I_i$, and consequently $x_1 x_2 \cdots x_{\rho_i} e_i \in I_i e_i$. On the other hand, $\mathcal{L}$ is a squarefree lexicographic submodule of F , then for all $u \in \{x_1, \dots, x_n\}^{\rho_i + f_i - f_{i-1}}$, we have that $ue_{i-1} >_{\text{slex}_F} x_1 x_2 \cdots x_{\rho_i} e_i$. Hence, $ue_{i-1} \in I_{i-1} e_{i-1}$, i.e., $u \in I_{i-1}$.

Conversely, let $\mathcal{L}$ be a graded submodule of F satisfying (i) and (ii). Since every ideal I_i is a squarefree lexsegment ideal, we have only to prove that for any pair (i, j) of integers with $1 \leq i < j \leq r$, if $ue_i, ve_j \in M^s(\mathcal{L}_d)$, then $ve_j \in \mathcal{L}$ implies $ue_i \in \mathcal{L}$, where $M^s(\mathcal{L}_d)$ is the set of all squarefree monomials of degree d of $\mathcal{L}$.

(Case 1). $i = j - 1$. Let $ue_{j-1}, ve_j \in M^s(\mathcal{L}_d)$ with $ve_j \in \mathcal{L}$.

Since $d = \deg ue_{j-1} = \deg ve_j$, it follows that $\deg u = \deg v + f_j - f_{j-1} \geq \rho_j + f_j - f_{j-1}$ and so $u \in [x_1, \dots, x_n]^{\rho_j + f_j - f_{j-1}} \subseteq I_{j-1}$.

(Case 2). $i \leq j - 2$. Let $ue_i >_{\text{slex}_F} ve_j$ with $ue_i, ve_j \in M^s(\mathcal{L}_d)$ and $ve_j \in \mathcal{L}$.

For $t = i + 1, \dots, j - 1$, set $w_t = x_1 x_2 \cdots x_{d-f_t}$.

It is

$$ue_i >_{\text{slex}_F} w_{i+1}e_{i+1} >_{\text{slex}_F} w_{i+2}e_{i+2} >_{\text{slex}_F} \cdots >_{\text{slex}_F} w_{j-1}e_{j-1} >_{\text{slex}_F} ve_j.$$

Since $d = \deg ue_i = \deg w_t e_t = \deg ve_j$, for $t = i + 1, \dots, j - 1$, then, from (Case 1), $w_{j-1} \in I_{j-1}$, $w_{j-2} \in I_{j-2}$, $\dots$, $w_{i+1} \in I_{i+1}$ and finally $u \in I_i$. $\square$

Example 2.9. (1) Let $S = K[x_1, x_2, x_3, x_4]$ and $F \simeq S^2$, $e_1 = (1, 0)$ and $e_2 = (0, 1)$. The submodule of F

$$\mathcal{L} = (x_1 x_2, x_1 x_3)e_1 \oplus (x_1 x_2 x_3, x_1 x_2 x_4)e_2$$

is not a squarefree lexicographic submodule of F even if the ideals $(x_1 x_2, x_1 x_3)$, $(x_1 x_2 x_3, x_1 x_2 x_4)$ are squarefree lexsegment ideals of S . In fact, $x_1 x_2 x_3 e_2 \in \mathcal{L}_3$ but $x_2 x_3 x_4 e_1 >_{\text{slex}_F} x_1 x_2 x_3 e_2$ and $x_2 x_3 x_4 e_1 \notin \mathcal{L}_3$. Observe that $[x_1, x_2, x_3, x_4]^3 \not\subseteq (x_1 x_2, x_1 x_3)$.

(2) Let $S = K[x_1, x_2, x_3, x_4, x_5]$ and $F \simeq S^3$, $e_1 = (1, 0, 0)$, $e_2 = (0, 1, 0)$ and $e_3 = (0, 0, 1)$. The submodule of F

$$\begin{aligned} \mathcal{L} = [x_1, x_2, x_3, x_4, x_5]^2 e_1 \oplus (x_1 x_2 x_3, x_1 x_2 x_4, x_1 x_2 x_5, x_1 x_3 x_4, x_2 x_3 x_4 x_5) e_2 \oplus \\ \oplus (x_1 x_2 x_3 x_4, x_1 x_2 x_3 x_5, x_1 x_2 x_4 x_5) e_3 \end{aligned}$$

is a squarefree lexicographic submodule of F .

Corollary 2.10. Let $\mathcal{L} = \bigoplus_{i=1}^r I_i e_i \subsetneq F$ be a squarefree lexicographic submodule.

Set

$$\mu_i = \max\{\deg ue_i : ue_i \in G(I_i e_i)\},$$

for $i = 1, \dots, r$. Then

- 1) $\mu_i - f_i \leq \text{indeg} I_{i+1} + f_{i+1} - f_i$, for $i = 1, \dots, r - 1$.
- 2) $\mu_1 \leq \mu_2 \leq \cdots \leq \mu_r$.

Proof. It is a consequence of the fact that if $\mathcal{L}$ is a squarefree lexicographic submodule then $[x_1, \dots, x_n]^{\rho_i + f_i - f_{i-1}} \subseteq I_{i-1}$, for $i = 2, \dots, r$, with $\rho_i = \text{indeg}(I_i)$. $\square$

The next result shows the relation between the class of squarefree lexicographic submodules and the class of squarefree (strongly) stable submodule.

Proposition 2.11. Let $\mathcal{L} = \bigoplus_{i=1}^r I_i e_i \subsetneq F$ be a squarefree lexicographic submodule, then $\mathcal{L}$ is a squarefree strongly stable submodule.

Proof. Since every squarefree lexsegment ideal is a squarefree strongly stable ideal [1], from Proposition 2.3, it is sufficient to show that

$$(2.2) \quad [x_1, \dots, x_n]^{f_{i+1}-f_i} I_{i+1} \subseteq I_i$$

for $i = 1, \dots, r-1$.

Since

$$[x_1, \dots, x_n]^{f_{i+1}-f_i} I_{i+1} \subseteq [x_1, \dots, x_n]^{f_{i+1}-f_i+\text{indeg} I_{i+1}},$$

the assert follows from Proposition 2.8. $\square$

Hence, we have the following hierarchy of submodules:
squarefree lexicographic submodule $\Rightarrow$ squarefree strongly stable submodule $\Rightarrow$ squarefree stable submodule.

3. EXTREMAL BETTI NUMBERS OF SQUAREFREE STABLE SUBMODULES

In this Section we examine the extremal Betti numbers of squarefree stable submodules.

If I is a squarefree stable ideal, then [1]:

$$(3.1) \quad \beta_{i,i+j}(I) = \sum_{u \in G(I)_j} \binom{m(u)-j}{i}.$$

Hence, since every squarefree stable submodule $M \subsetneq F$ is a squarefree monomial submodule, we have that

$$(3.2) \quad \beta_{k,k+j}(M) = \sum_{i=1}^r \beta_{k,k+j-f_i}(I_i) = \sum_{i=1}^r \left[\sum_{u \in G(I_i)_{j-f_i}} \binom{m(u)-j+f_i}{k} \right].$$

The next result shows that all linear strands of a squarefree stable submodule begin in homological degree 0.

For a positive integer t , we set $[t] = \{1, \dots, t\}$.

Lemma 3.1. *Let $M = \bigoplus_{t=1}^r I_t e_t \subsetneq F$ be a squarefree stable submodule. If $\beta_{i,i+j}(M) \neq 0$, then $\beta_{k,k+j}(M) \neq 0$ for $k = 0, \dots, i$.*

Proof. If $\beta_{i,i+j}(M) \neq 0$, by (3.2) there exists $t \in [r]$ and $u \in G(I_t)_{j-f_t}$ such that $m(u) \geq i+j-f_t$.

It follows that $m(u) \geq k+j-f_t$, for $k = 0, \dots, i$. Hence, $\beta_{k,k+j-f_t}(I_t) \neq 0$ and $\beta_{k,k+j}(M) \neq 0$, for $k = 0, \dots, i$. $\square$

From Definition 1.1, it follows:

Corollary 3.2. *Under the same hypotheses of Lemma 3.1 for M . The following conditions are equivalent:*

- (a) $\beta_{k,k+\ell}(M)$ is extremal;
- (b) (1) $\beta_{k,k+\ell}(M) \neq 0$;
(2) $\beta_{k,k+j}(M) = 0$, for $j > \ell$;
(3) $\beta_{i,i+\ell}(M) = 0$, for $i > k$.

Theorem 3.3. *Let $M = \oplus_{i=1}^r I_i e_i \subsetneq F$ be a squarefree stable submodule.*

The following conditions are equivalent:

- (1) $\beta_{k,k+\ell}(M)$ is extremal;
- (2) $k + \ell = \max\{m(u) + f_i : ue_i \in G(M)_\ell, i = 1, \dots, r\}$ and $m(u) + f_i < k + j$, for all $j > \ell$ and for all $ue_i \in G(M)_j$.

Proof. (1) $\Rightarrow$ (2). By (3.2) $\beta_{k,k+\ell}(M) \neq 0$ if and only if there exists a squarefree monomial $ue_i \in G(M)_\ell$ such that $m(u) + f_i \geq k + \ell$, for some $i \in [r]$. As a consequence, $\max\{m(u) + f_i : ue_i \in G(M)_\ell, i = 1, \dots, r\} \geq k + \ell$.

Suppose $t + \ell := \max\{m(u) + f_i : ue_i \in G(M)_\ell, i = 1, \dots, r\} > k + \ell$. Hence, $\beta_{t,t+\ell}(M) \neq 0$, for $t > k$. This is a contradiction from Corollary 3.2, (b), whereupon

$$k + \ell = \max\{m(u) + f_i : ue_i \in G(M)_\ell, i = 1, \dots, r\}.$$

Suppose there exists $j > \ell$ and a squarefree monomial $ue_i \in G(M)_j$, for some $i \in [r]$, such that $m(u) + f_i \geq k + j$. From (3.2), then $\beta_{k,k+j}(M) \neq 0$. Again a contradiction from Corollary 3.2, (b).

(2) $\Rightarrow$ (1). Since $k + \ell = \max\{m(u) + f_i : ue_i \in G(M)_\ell, i = 1, \dots, r\}$, then $\beta_{k,k+\ell}(M) \neq 0$ and $\beta_{i,i+\ell}(M) = 0$, for all $i > k$. On the other hand, $m(u) + f_i < k + j$, for all $j > \ell$ and for all $ue_i \in G(M)_j$, implies $\beta_{k,k+j}(M) = 0$. Therefore, from Corollary 3.2, we get the assert. $\square$

As consequences, we obtain the following corollaries.

Corollary 3.4. *Let $M = \oplus_{i=1}^r I_i e_i \subsetneq F$ be a squarefree stable submodule and let $\beta_{k,k+\ell}(M)$ an extremal Betti number of M . Then*

$$\beta_{k,k+\ell}(M) = |\{ue_i \in G(M)_\ell : m(u) + f_i = k + \ell, i = 1, \dots, r\}|.$$

Corollary 3.5. *Let $M = \oplus_{i=1}^r I_i e_i \subsetneq F$ be a squarefree stable submodule.*

Set

$$\ell = \max\{j : G(M)_j \neq \emptyset\}$$

and

$$m = \max\{m(u) + f_i : ue_i \in G(M), i = 1, \dots, r\}.$$

Then $\beta_{m-\ell,m}$ is the unique extremal Betti number of M if and only if

$$m = \max\{m(u) + f_i : ue_i \in G(M)_\ell, i = 1, \dots, r\},$$

and for every $w \in G(M)_j, j < \ell, m(w) < m$.

Remark 3.6. *Under the same hypotheses of Theorem 3.3, if $F \simeq S^r$, $\beta_{k,k+\ell}(M)$ is extremal if and only if*

$$k + \ell = \max\{m(u) : ue_i \in G(M)_\ell, i = 1, \dots, r\},$$

and $m(u) < k + j$, for all $j > \ell$ and for all $ue_i \in G(M)_j$.

Moreover,

$$\beta_{k,k+\ell}(M) = |\{ue_i \in G(M)_\ell : m(u) = k + \ell, i = 1, \dots, r\}|.$$

Remark 3.7. If I is a squarefree stable monomial ideal of S and $\beta_{k,k+\ell}(I)$ is an extremal Betti number of I , then from (3.2) and Remark 3.6, we have

$$(3.3) \quad 1 \leq \beta_{k,k+\ell}(I) \leq \binom{k+\ell-1}{\ell-1}.$$

In fact, there are exactly $\binom{k+\ell-1}{\ell-1}$ squarefree monomials of degree ℓ in S with $m(u) = k + \ell$.

If $M = \oplus_{i=1}^r I_i e_i \subsetneq F$ is a squarefree stable submodule and $\beta_{k,k+\ell}(M)$ is an extremal Betti number of M , then there exist some squarefree stable ideals $I_{j_1}, \dots, I_{j_t}$, $\{j_1, \dots, j_t\} \subseteq [n]$, $1 \leq j_1 < j_2 < \dots < j_t \leq r$, with $(k, \ell - f_{j_1}), \dots, (k, \ell - f_{j_t})$ as corners.

Then

$$\beta_{k,k+\ell}(M) = \sum_{i=1}^t \beta_{k,k+\ell}(I_{j_i} e_{j_i}) = \sum_{i=1}^t \beta_{k,k+\ell-f_{j_i}}(I_{j_i}),$$

and

$$1 \leq \beta_{k,k+\ell}(M) \leq \sum_{i=1}^t \binom{k+\ell-f_{j_i}-1}{\ell-f_{j_i}-1},$$

where $\binom{k+\ell-f_{j_i}-1}{\ell-f_{j_i}-1}$ is the number of all squarefree monomials u of S of degree $\ell - f_{j_i}$ with $m(u) = k + \ell$.

4. A CRITERION FOR DETERMINING EXTREMAL BETTI NUMBERS

In this Section, we describe a criterion for determining the extremal Betti numbers of a squarefree stable submodule of the graded free S -module S^r , $r \geq 1$.

Let $M = \oplus_{i=1}^r I_i e_i \subsetneq S^r$ be a squarefree stable submodule.

If M is generated in one degree d , then M has a unique extremal Betti number $\beta_{m-d,m}(M)$, where $m = \max\{m(u) : ue_i \in G(M), i = 1, \dots, r\}$.

Let M be generated in degrees $1 \leq d_1 < d_2 < \dots < d_t \leq n$.

Set

$$m_{d_j} = \max\{m(u) : ue_i \in G(M)_{d_j}, i = 1, \dots, r\},$$

for $j = 1, \dots, t$.

Consider the following sequence of non negative integers associated to M :

$$(4.1) \quad \mathbf{ds}(M) = (m_{d_1} - d_1, m_{d_2} - d_2, \dots, m_{d_t} - d_t).$$

We call it the *degree-sequence* of M .

Remark 4.1. Assume that for some $j \in [t]$, $G(M)_{d_j} = \{x_1 \cdots x_{d_j}\}$. Then $m_{d_j} - d_j = 0$.

If $F = S$, and consequently M is a squarefree monomial ideal in S , then $m_{d_i} - d_i > 0$, for $i = 2, \dots, t$. Moreover, if $G(M)_{d_1} = G(M)_{\text{indeg}(M)} = \{x_1 \cdots x_{\text{indeg}(M)}\}$, then $m_{d_1} - d_1 = 0$.

We can observe that, if

$$(4.2) \quad m_{d_1} - d_1 > m_{d_2} - d_2 > \dots > m_{d_t} - d_t,$$

then, from Theorem 3.3, for $i = 1, \dots, t$, $\beta_{m_{d_i}-d_i, m_{d_i}}(M)$ is an extremal Betti number of M .

Suppose that (4.2) does not hold.

We construct a suitable subsequence of the *degree-sequence* $\mathbf{ds}(M)$:

$$(4.3) \quad \widehat{\mathbf{ds}}(M) = (m_{d_{i_1}} - d_{i_1}, m_{d_{i_2}} - d_{i_2}, \dots, m_{d_{i_q}} - d_{i_q}),$$

with $d_1 \leq d_{i_1} < d_{i_2} < \dots < d_{i_q} = d_t$ and such that, for $j = 1, \dots, q$, $\beta_{m_{d_{i_j}}-d_{i_j}, m_{d_{i_j}}}(M)$ is an extremal Betti number of M .

Construction 4.2. The admissible subsequence $\widehat{\mathbf{ds}}(M)$ is obtained as follows:

d_{i_1} is the smallest degree of a monomial generator of M such that

$$m_{d_{i_1}} - d_{i_1} > m_{d_j} - d_j, \text{ for all } j > i_1;$$

$d_{i_2} > d_{i_1}$ is the smallest degree of a monomial generator of M such that

$$m_{d_{i_2}} - d_{i_2} > m_{d_j} - d_j, \text{ for all } j > i_2 > i_1;$$

.....

$d_{i_{q-1}} > d_{i_{q-2}}$ is the smallest degree of a monomial generator of M such that

$$m_{d_{i_{q-1}}} - d_{i_{q-1}} > m_{d_j} - d_j, \text{ for all } j > i_{q-1} > \dots > i_1;$$

$$d_{i_q} = d_t.$$

It is $m_{d_{i_1}} - d_{i_1} > m_{d_{i_2}} - d_{i_2} > \dots > m_{d_{i_q}} - d_{i_q}$, and Theorem 3.3 guarantees that

$$\beta_{m_{d_{i_1}}-d_{i_1}, m_{d_{i_1}}}(M), \beta_{m_{d_{i_2}}-d_{i_2}, m_{d_{i_2}}}(M), \dots, \beta_{m_{d_{i_q}}-d_{i_q}, m_{d_{i_q}}}(M)$$

are extremal Betti numbers of M .

The integer q is called the *degree-length* of M and gives the number of the extremal Betti numbers of the squarefree stable submodule M . We will denote such integer by $\mathbf{dl}(M)$.

Examples 4.3. (1) Let $S = K[x_1, x_2, x_3, x_4, x_5, x_6, x_7]$ and let

$$I = (x_1x_2, x_1x_3, x_1x_4, x_1x_5, x_1x_6, x_2x_3x_4, x_2x_3x_5, x_2x_4x_5, x_2x_3x_6x_7, x_3x_4x_5x_6x_7)$$

be a squarefree strongly stable ideal of S .

The *degree-sequence* of I is

$$\mathbf{ds}(I) = (m_2 - 2, m_3 - 3, m_4 - 4, m_5 - 5) = (4, 2, 3, 2).$$

Following Construction 4.2, the admissible subsequence $\widehat{\mathbf{ds}}(I)$ is $(4, 3, 2)$ and $\mathbf{dl}(I) = 3$. The extremal Betti numbers of I are

$$\beta_{6-2,6}(I), \beta_{7-4,7}(I), \beta_{7-5,7}(I),$$

as the Betti table of I shows:

		0	1	2	3	4
2	:	5	10	10	5	1
3	:	3	5	2	—	—
4	:	1	3	3	1	—
5	:	1	2	1	—	—

(2) Let $S = K[x_1, x_2, x_3, x_4, x_5, x_6]$ and $F \simeq S^4$, $e_1 = (1, 0, 0, 0)$, $e_2 = (0, 1, 0, 0)$, $e_3 = (0, 0, 1, 0)$, $e_4 = (0, 0, 0, 1)$. Let

$$M = I_1 e_1 \oplus I_2 e_2 \oplus I_3 e_3 \oplus I_4 e_4$$

with

$$I_1 = (x_1 x_2, x_1 x_3, x_1 x_4, x_2 x_3),$$

$$I_2 = (x_1 x_2 x_3, x_1 x_2 x_4, x_1 x_3 x_4, x_1 x_3 x_5, x_1 x_3 x_6, x_2 x_3 x_4, x_2 x_3 x_5, x_2 x_3 x_6),$$

$$I_3 = (x_1 x_2 x_3 x_4, x_1 x_2 x_3 x_5, x_1 x_2 x_4 x_5, x_1 x_2 x_4 x_6, x_2 x_3 x_4 x_5), I_4 = (x_1 x_2 x_3 x_4 x_5),$$

be a squarefree stable submodule of F .

The *degree-sequence* of M is

$$\mathbf{ds}(M) = (m_2 - 2, m_3 - 3, m_4 - 4, m_5 - 5) = (2, 3, 2, 0).$$

Following Construction 4.2, the admissible subsequence $\widehat{\mathbf{ds}}(M)$ is $(3, 2, 0)$ and $\mathbf{dl}(M) = 3$. Indeed, the extremal Betti numbers of M are

$$\beta_{6-3,6}(M), \beta_{6-4,6}(M), \beta_{5-5,5}(M),$$

as one can read on its Betti table:

		0	1	2	3
2	:	4	4	1	—
3	:	8	13	8	2
4	:	5	5	1	—
5	:	1	—	—	—

5. AN APPLICATION

In this Section, we consider some special extremal Betti numbers of squarefree lexicographic submodules.

If M is a finitely generated graded $S = K[x_1, \dots, x_n]$ -module, then $\beta_{i,j}(M) = 0$, for all i and $j > n$. Therefore, $\beta_{i,n}(M)$ is an extremal Betti number if $\beta_{i,n}(M) \neq 0$. Such extremal Betti numbers are called *super extremal* [2].

The pair $(i, n - i)$ is called a *super corner*.

Let $M = \bigoplus_{i=1}^r I_i e_i \subsetneq S^r$, $r \geq 1$, be a squarefree stable submodule and $G(M)$ its minimal system of monomial generators.

Set

$$b_i = |\{ue_j \in G(M)_{n-i} : m(u) = n, j = 1, \dots, r\}|,$$

then $\beta_{i,n}(M) = b_i$, for $i = 1, \dots, n - 1$. We call

$$\mathbf{b}(M) = (b_0, b_1, \dots, b_{n-1})$$

the $\mathbf{b}$ -vector of M .

For $\mathbf{b}(I) = (b_0, b_1, \dots, b_{n-1})$, we define the *support* of $\mathbf{b}(M)$ to be the following set:

$$\text{supp}(\mathbf{b}(M)) = \{i \in \{0, 1, \dots, n-1\} : b_i \neq 0\}.$$

For every $u \in \{x_1, \dots, x_n\}^d$, we denote by $\langle u \rangle$ the squarefree lexsegment ideal of degree d in S defined as follows:

$$\langle u \rangle = (w \in \{x_1, \dots, x_n\}^d : w \geq_{\text{slex}} u).$$

Lemma 5.1. *Let $n \geq 3$ and X a non-empty subset of $\{0, 1, \dots, n-1\}$.*

For every integer $r \geq |X|$ there exists a squarefree lexicographic submodule $\mathcal{L} \subsetneq S^r$ with $\text{supp}(\mathbf{b}(\mathcal{L})) = X$.

Proof. Set $X = \{k_1, k_2, \dots, k_t\}$ with $k_1 < k_2 < \dots < k_t$ and $r \geq t$.

By Proposition 2.8, an admissible squarefree lexicographic submodule $\mathcal{L} \subsetneq S^r$ such that $\text{supp}(\mathbf{b}(\mathcal{L})) = X$ is:

$$\mathcal{L} = \bigoplus_{j=0}^{t-2} [x_1, \dots, x_n]^{n-k_{t-j}} e_{j+1} \oplus \left(\bigoplus_{i=t}^{r-1} [x_1, \dots, x_n]^{n-k_1} e_i \right) \oplus \langle x_1 x_2 \cdots x_{n-k_1-1} x_n \rangle e_r.$$

Note that $\mathbf{b}(\mathcal{L}) = (b_0, \dots, b_{n-1})$, where $b_i = 0$, for $i \in \{0, 1, \dots, n-1\} \setminus X$, $b_{k_j} = \binom{n-1}{n-k_j-1}$, for $j = 2, \dots, t$ and $b_{k_1} = 1 + (r-t) \binom{n-1}{n-k_1-1}$. □

Proposition 5.2. *Given three integers n, t, r such that $n \geq 3$, $1 \leq t \leq n-1$, $r \geq t$, t pairs of integers $(k_1, \ell_1), (k_2, \ell_2), \dots, (k_t, \ell_t)$, with*

$$0 \leq k_t < k_{t-1} < \dots < k_1 \leq n-1, \quad 1 \leq \ell_1 < \ell_2 < \dots < \ell_t \leq n$$

and such that, for $i = 1, \dots, t$, $k_i + \ell_i = n$.

Then there exists a squarefree lexicographic submodule $\mathcal{L} \subsetneq S^r$, generated in degrees $\ell_1, \ell_2, \dots, \ell_t$ with $(k_1, \ell_1), (k_2, \ell_2), \dots, (k_t, \ell_t)$ as super corners.

Proof. Set $X = \{n-\ell_t, n-\ell_{t-1}, \dots, n-\ell_1\} = \{k_t, k_{t-1}, \dots, k_1\}$. One has that X is a non-empty subset of $\{0, 1, \dots, n-1\}$ and from Lemma 5.1 the squarefree monomial submodule of S^r :

$$\mathcal{L} = \bigoplus_{i=1}^{t-1} [x_1, \dots, x_n]^{\ell_i} e_i \oplus \left(\bigoplus_{i=t+1}^{r-1} [x_1, \dots, x_n]^{\ell_t} e_i \right) \oplus \langle x_1 x_2 \cdots x_{\ell_t-1} x_n \rangle e_r$$

is a squarefree lexicographic submodule generated in degrees $\ell_1, \ell_2, \dots, \ell_t$ and such that $\text{supp}(\mathbf{b}(\mathcal{L})) = X$. Thus, $b_{k_i} = \beta_{k_i, n}$ are super extremal Betti numbers of $\mathcal{L}$, and the assert follows. □

Remark 5.3. *Let $n = 2$. If one considers the pair $(k, \ell) = (0, 2)$, then there exists the squarefree lexicographic submodule $\mathcal{L} = \bigoplus_{i=1}^r (x_1 x_2) e_i$ with $\mathbf{b}(\mathcal{L}) = (r, 0)$.*

If one considers $(k, \ell) = (1, 1)$, then there exists the squarefree lexicographic submodule $\mathcal{L} = \bigoplus_{i=1}^r (x_1, x_2) e_i$ with $\mathbf{b}(\mathcal{L}) = (0, r)$.

REFERENCES

- [1] A. Aramova, J. Herzog and T. Hibi, Squarefree lexsegment ideals, *Math.Z.* **228** (1998), 353–378.
- [2] A. Aramova, J. Herzog and T. Hibi, Shifting operations and graded Betti numbers ideals, *J. Algebraic Combin.* **12** (2000), 207–222.
- [3] D. Bayer, H. Charalambous, and S. Popescu. Extremal Betti numbers and Applications to Monomial Ideals, *J. Algebra* **221** (1999), 497–512.
- [4] W. Bruns, J. Herzog. *Cohen-Macaulay rings* **39** Revised Edition, Cambridge 1996.
- [5] A. Capani, A. Niesi, L. Robbiano, *CoCoA: A system for doing computations in Commutative Algebra*, available at <http://cocoa.dima.unige.it>.
- [6] M. Crupi, M. La Barbiera. Algebraic properties of universal squarefree lexsegment ideals, *Algebra Colloq.*, to appear (2013).
- [7] M. Crupi, R. Utano. Extremal Betti numbers of lexsegment ideals, *Lecture notes in Pure and Applied Math., Geometric and combinatorial aspects of Commutative algebra*, **217**, (2000), 159–164.
- [8] M. Crupi, R. Utano. Extremal Betti numbers of graded ideals, *Results Math.* **43**, (2003), 235–244.
- [9] M. Crupi, R. Utano. Minimal resolutions of some monomial submodules, *Results Math.* **55**, (2009), 311–328.
- [10] M. Crupi, G. Restuccia. Monomial modules and graded Betti numbers, *Math. Notes* **85**, (2009), 690–702.
- [11] S. Eliahou, M. Kervaire. Minimal resolutions of some monomial ideals, *J. Algebra*, **129** (1990), 1–25.
- [12] D. Eisenbud, *Commutative Algebra with a view toward Algebraic Geometry*, Springer-Verlag, 1995.
- [13] D. R. Grayson and M.E. Stillman, *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www.math.uiuc.edu/Macaulay2>.
- [14] K. Pardue. Deformation classes of graded modules and maximal Betti numbers Illinois *J. Math.* **40**, N. 4 (1996), 564–585.